

Instrumental developments at the Gregory-Coudé Telescope at IRSOL

M. Bianda^{1,2,*}, R. Ramelli¹, A. Feller², J.O. Stenflo^{2,3}, and G. Küveler⁴

¹Istituto Ricerche Solari Locarno, Locarno, Switzerland

²Institute of Astronomy, ETH Zentrum, Zurich, Switzerland

³Faculty of Mathematics & Science, University of Zurich, Zurich, Switzerland

⁴Fachhochschule Wiesbaden, Rüsselsheim, Germany

**Email:* mbianda@irsol.ch

Abstract. The research projects carried out with the Gregory-Coudé Telescope (GCT) at Istituto Ricerche Solari Locarno (IRSOL) are mainly focused on high precision polarimetry. The Zurich IMaging POLarimeter (ZIMPOL) developed at ETH Zurich and installed permanently at the GCT at IRSOL allows a polarimetric precision down to 10^{-5} to be reached. This makes it possible to perform several accurate spectro-polarimetric measurements of scattering polarization and to investigate solar magnetic fields through the signatures of the Hanle and Zeeman effects. The research programs are currently being extended to monochromatic imaging of the Stokes vector with a recently installed Fabry-Perot rapidly tunable filter system with a narrow pass band of about 30 mÅ. The spatial resolution is being improved by the installation of an adaptive optics (AO) system.

1 Introduction

The great advances in high precision polarimetry that have been achieved with the introduction of the Zurich IMaging POLarimeter (ZIMPOL) a decade ago opened a new window in solar physics. Polarimetry is in fact a very powerful tool that can be used to study solar magnetic fields as well as the physical processes behind the generation of polarization in atomic and molecular spectral lines. Magnetic field measurements through Zeeman effect signatures, which appear in the presence of strong and oriented magnetic fields, have long been performed at many observatories. With the high polarimetric precision of ZIMPOL it has become possible to extend the magnetic field diagnostics to weak fields and to fields which are tangled on scales below the spatial resolution, which are invisible to the Zeeman effect but get revealed by the Hanle effect (Hanle 1924).

Spectro-polarimetry is currently the main field of research at the Istituto Ricerche Solari Locarno (IRSOL). Advantage is taken from the circumstance that a ZIMPOL system is permanently installed at IRSOL. In addition, the Gregory-Coudé Telescope of the observatory is very well suited for polarimetric measurements, since the amount of instrumental polarization is low and stays practically constant during the observing day, since it is a function of declination only. Therefore it can easily be accounted for. With a Fabry-Perot filter system and an adaptive optics system recently installed at IRSOL we plan to start several new interesting projects.

2 IRSOL - The institute

The observatory at the Istituto Ricerche Solari Locarno (IRSOL), located in southern Switzerland, was constructed in 1958/59 by the Universitäts-Sternwarte Göttingen (USG), Germany. In 1984, after USG moved its observing activity to the facilities at Observatorio del Teide on Tenerife, a local foundation (FIRSOL) acquired the observatory in Locarno. The partially dismantled instrumentation was rebuilt and improved, in collaboration with USG (now IAG), Technical Highschool (FH) of Applied Sciences of Wiesbaden (Germany), and the Institute of Astronomy at ETH Zurich. The scientific collaboration with ETH Zurich allowed the implementation at IRSOL of an important polarimetry observing program, first with a beam exchange polarimeter and then with ZIMPOL.

3 Instrumentation at IRSOL

The IRSOL telescope is a 45 cm aperture Gregory coudé type instrument with 25 m effective focal length. A field stop with a pinhole of about 2.5 mm diameter in the prime focus reduces the field of view to a 200 arcsec diameter circular image. The rest of the solar image is reflected away from the main light beam. This reduces heating and scattered light and is of particular advantage when observing low intensity structures like sunspots, spicules, or prominences. The relative orientation of the two folding mirrors M3 and M4 (coudé) changes only with declination and is orthogonal at the time of the equinoxes. As a consequence the instrumental polarization, originating through oblique reflections, is almost constant during the day and virtually vanishes during the equinoxes (Sánchez Almeida et al. 1991). A Gregory-Coudé telescope is thus very well suited for polarimetric measurements.

An automatic guiding system developed by the FH Wiesbaden (Küveler et al. 1998) is also available. Its operation is based on the solar image obtained from the light reflected by the field stop in the primary focus.

The Czerny-Turner spectrograph with 10 m focal length is based on a 180×360 mm grating with 316 lines per mm and 63° Blaze angle. A prism based predisperser allows to select the spectral band entering in the spectrograph without overlap of the grating orders.

Monochromatic imaging observations of the solar surface can be performed using the recently installed Fabry-Perot filter system in collimated configuration (Feller et al. 2006). The system uses two temperature controlled lithium niobate etalons with an aperture of 70 mm. The transmitted wavelength can be selected by electrically tuning the refractive index of the etalon medium, by varying the temperature, or by tilting the etalon. The bandwidth is about 30 mÅ.

An adaptive optics (AO) system based on a tip-tilt mirror and a 37 actuator deformable mirror is currently being installed and tested in collaboration with the University of Applied Sciences of Southern Switzerland, SUPSI, and with ETH Zurich. The system follows the design of the infrared AO system installed at the McMath-Pierce Solar Telescope at Kitt Peak (Keller et al. 2003). The first tests made with the tip-tilt mirror have already given good results (see Figure 1).

Two polarimeters are available. The oldest one operating at IRSOL is the dual beam exchange device based on a Savart plate and rotating quarter and half wave retarder plates (Bianda et al. 1998). A polarimetric precision of a few 10^{-4} can be reached, but it can be affected by seeing-induced cross-talk, because the technique requires two exposures taken




Figure 1. First tests with the tip-tilt correction of the AO system. The two plots show the displacement of the barycentre of a sunspot in the x resp. y direction with and without tip-tilt correction. The tip-tilt mirror was switched off around frame number 1200.

at different times. The second polarimeter is ZIMPOL (Povel 1995; Gandorfer et al. 2004), which is installed permanently at IRSOL since 1998. Its main advantage is that it is free from seeing-induced effects thanks to its high modulation rate: 42 kHz (obtained with a piezoelectric modulator), or 1 kHz (obtained with ferro-electric liquid crystal modulators). Another advantage is that the same pixels of the CCD ZIMPOL sensor are used to measure all Stokes parameters. Therefore the Stokes Q/I , U/I and V/I images are not influenced by different pixel efficiencies. The ZIMPOL polarimetric accuracy depends mainly on the photon statistics. With long exposure times it has already been possible at IRSOL to reach an accuracy of about 10^{-5} .

4 Scientific research projects at IRSOL

The research projects at IRSOL take advantage of the very good polarimetric and spectral accuracy of the instrumentation. A large amount of observing time is available to carry out monitoring measurements or for projects requiring long observing times (which cannot easily be done at large telescope facilities, where the observing time is shared by different research groups according to a predefined program). In addition it is possible to be very flexible with the programs to allow fast reaction to particular solar events. The modular layout of the instrumentation inside the observing room is very convenient for installation and testing of new instrumentation.

Examples of the scientific results obtained at IRSOL in recent years are:

- Investigations of the Hanle effect in the quiet chromosphere (Bianda et al. 1999).
- Publication of the first two volumes of the “Atlas of the Second Solar Spectrum” (Gandorfer 2000, 2002).
- Discovery of vast amounts of hidden magnetism in the solar photosphere (Trujillo Bueno et al. 2004; Stenflo 2004).
- Determination of novel constraints on impact polarization in solar flares (Bianda et al. 2005).
- Measurements of full Stokes profiles in prominences in H_{α} , He D₃, H_{β} , and in spicules in He D₃ (Ramelli et al. 2005a,b).
- First polarimetric measurements of the Paschen-Back effect in CaH transitions (Berdyugina et al. 2006).

Different other observing programs are also foreseen in the future. They will focus on solar magnetism and polarimetry with the Fabry-Perot filter system or with the spectrograph. They also include synoptic type programs (e.g. variations of the Hanle-effect signatures with heliographic latitude and solar cycle). Furthermore IRSOL is open to coordinated type programs with other observatories: simultaneous observations of solar features with complementary sets of instruments or supporting type observations that complement the science of another project.

Acknowledgements. We are grateful for the financial support provided by the canton of Ticino, the city of Locarno, ETH Zurich and the Fondazione Aldo e Cele Daccò.

References

- Berdyugina, S. V., Fluri, D. M., Ramelli, R., Bianda, M., Gisler, D., and Stenflo, J. O. 2006, *ApJ*, **649**, L49
- Bianda, M., Solanki, S. K., and Stenflo, J. O. 1998, *A&A*, **331**, 760
- Bianda, M., Stenflo, J. O., and Solanki, S. K. 1999, *A&A*, **350**, 1060
- Bianda, M., Benz, A. O., Stenflo, J. O., Küveler, G., Ramelli, R. 2005, *A&A*, **434**, 1183
- Feller, A., et al. 2006, Proceedings of the Fourth Solar Polarization Workshop, Boulder 2005
- Gandorfer, A. 2000, VdF, ISBN 3-7281-2764-7
- Gandorfer, A. 2002, VdF, ISBN 3-7281-2844-4
- Gandorfer, A., et al. 2004, *A&A*, **422**, 703
- Hanle, W. 1924, *Z. Phys.*, **30**, 93
- Keller, C. U., Plymate, C. and Ammons, S. M. 2003, Proceedings of the SPIE, **4853**, 35
- Küveler, G., et al. 1998, *Solar Phys.*, **182**, 247
- Povel, H. P. 1995, *Opt. Eng.* **34**, 1870
- Ramelli, R., and Bianda, M. 2005a, *Astron. Astrophys. Space Science Library*, **320**, 215
- Ramelli, R., Bianda, M., Trujillo Bueno, J., Merenda, L., and Stenflo, J. O. 2005b, in D. E. Innes, A. Lagg, and S. A. Solanki (Eds.), *ESA SP-596: Chromospheric and Coronal Magnetic Fields*
- Sánchez Almeida, J., Martínez Pillet, V., and Wittmann, A. D. 1991, *Solar Phys.* **134**, 1
- Stenflo, J. O. 2004, *Nature*, **430**, 304
- Trujillo Bueno, J., Shchukina, N., and Asensio Ramos, A. 2004, *Nature*, **430**, 326